

Canonicity of Bäcklund transformation: r -matrix approach. II.

E K Sklyanin ¹

Laboratoire de Physique ², Groupe de Physique Théorique, ENS Lyon,
46 allée d'Italie, 69364 Lyon 07, France

Abstract. This is the second part of the paper devoted to the general proof of canonicity of Bäcklund transformation (BT) for a Hamiltonian integrable system governed by $SL(2)$ -invariant r -matrix. Introducing an extended phase space from which the original one is obtained by imposing a 1st kind constraint, we are able to prove the canonicity of BT in a new way. The new proof allows to explain naturally the fact why the gauge transformation matrix M associated to the BT has the same structure as the Lax operator L . The technique is illustrated on the example of the DST chain.

25 March, 1999

To be published in: *Trudy MIAN*, v. 226 (1999), Moscow, Nauka

¹On leave from: Steklov Mathematical Institute at St. Petersburg, Fontanka 27, St. Petersburg 191011, Russia. E-mail: sklyanin@euclid.pdmi.ras.ru

²UMR 5672 du CNRS et de l'ENS Lyon

1. Introduction

In the first part of this paper [1] we have addressed the problem of proving the canonicity of Bäcklund transformation for the Hamiltonian integrable models associated to $SL(2)$ -invariant r -matrices (such as Heisenberg magnet, Toda lattice, Nonlinear Schrödinger equation).

Consider a finite-dimensional Hamiltonian system defined in terms of canonical variables (X, x) :

$$\{X_i, X_j\} = \{x_i, x_j\} = 0, \quad \{X_i, x_j\} = \delta_{ij}. \quad (1.1)$$

Suppose that the system is completely integrable and possesses a Lax matrix $L(u; X, x)$ whose spectral invariants generate the commuting Hamiltonians H_n (u being a complex parameter). In what follows we often shall make use of several sets of canonical variables, like (X, x) , (Y, y) , (S, s) , (T, t) etc. To simplify the notation, we shall mark the Lax matrix expressed in terms of the corresponding variables as, for example, $L^{(x)}(u) \equiv L(u; X, x)$ omitting the subscript when it is not important.

As in [1], we assume that $L(u)$ is a matrix of order 2×2 and the Poisson brackets between the entries of $L(u)$ can be expressed in the r -matrix form [2]

$$\{\overset{1}{L}(u), \overset{2}{L}(v)\} = [r_{12}(u-v), \overset{1}{L}(u)\overset{2}{L}(v)] \quad (1.2)$$

where $\overset{1}{L} = L \otimes \mathbb{1}$, $\overset{2}{L} = \mathbb{1} \otimes L$, and $r_{12} = \kappa(u-v)^{-1}\mathcal{P}_{12}$ is the standard $SL(2)$ -invariant solution to the classical Yang-Baxter equation [2], κ being a constant and $\mathcal{P}_{12}$ being the permutation operator in $\mathbb{C}^2 \otimes \mathbb{C}^2$.

A Bäcklund transformation, by definition [3], is a one-parametric family $\mathcal{B}_\lambda$ of commuting $\mathcal{B}_\lambda \circ \mathcal{B}_\mu = \mathcal{B}_\mu \circ \mathcal{B}_\lambda \forall \lambda, \mu \in \mathbb{C}$ canonical transformations $\mathcal{B}_\lambda : (X, x) \rightarrow (Y, y)$ preserving the Hamiltonians H_n . Since the Hamiltonians, in our case, are the spectral invariants of $L(u)$ there must exist a matrix $M_\lambda(u)$ such that

$$M_\lambda(u)L^{(x)}(u) = L^{(y)}(u)M_\lambda(u). \quad (1.3)$$

Here it is assumed that $Y = Y(X, x)$ and $y = y(X, x)$, so $M_\lambda(u) = M_\lambda(u; X, x)$. Typically, $M_\lambda(u)$ depends on the difference $u - \lambda$ which we shall assume henceforth.

In [1] and in the present paper we study, so to say, inverse problem, namely, given an integrable system and a Lax matrix, find such a matrix $M(u)$ that the similarity transformation

$$M(u - \lambda)L^{(x)}(u) = L^{(y)}(u)M(u - \lambda) \quad (1.4)$$

induces a canonical transformation $\mathcal{B}_\lambda : (X, x) \rightarrow (Y, y)$.

In [1] we have proved by a direct calculation the canonicity of the two-parametric transformation $\mathcal{B}_{\lambda_1 \lambda_2}$ corresponding to the matrix $M_{\lambda_1 \lambda_2}(u)$ of the form

$$M_{\lambda_1 \lambda_2}(u) = \begin{pmatrix} u - \lambda_1 + pq & p \\ -pq^2 + 2\mu q & u - \lambda_2 - pq \end{pmatrix}. \quad (1.5)$$

The proof, though straightforward and simple, has left one question unanswered, namely: why the ansatz (1.5) for $M_{\lambda_1 \lambda_2}(u)$ coincides with the elementary Lax matrix

$L(u)$ for Heisenberg XXX magnet [2] having the Poisson brackets (1.2) provided the variables p and q are canonical (note that in $M_{\lambda_1\lambda_2}(u)$ the variables pq , as shown in [1], *are not* canonical but have rather complicated brackets).

In the present paper we resolve this mystery and show that the abovementioned similarity between $M(u)$ and $L(u)$ is by no means accidental but, on the contrary, quite natural. The idea is to introduce an extended phase space from which the original one is obtained by imposing a 1st kind constraint. Such an indirect approach turns out not only to be simpler than the straightforward method of [1] but allows also to construct in a systematic way multiparametric families of Bäcklund transformations.

In the next section we give the general construction and a proof of canonicity of Bäcklund transformations. In section 3 we modify our construction to the case of integrable chains and in section 4 illustrate it on the example of the so-called DST chain (a degeneration of isotropic Heisenberg magnet).

2. General scheme

The first step of our construction is to extend the phase space by supplementing the original canonical variables (X, x) with an independent set of auxiliary canonical variables (S, s) . Note that the variables (S, s) commute with (X, x) and the dimension of the space (S, s) has no relation to the dimension of (X, x) .

Let matrix $M^{(s)}(u)$ has the same Poisson brackets (1.2) as $L(u)$

$$\{\overset{1}{M}(u), \overset{2}{M}(v)\} = [r_{12}(u-v), \overset{1}{M}(u)\overset{2}{M}(v)]. \quad (2.1)$$

Suppose that there exists a canonical transformation $\mathcal{R}_\lambda : (X, x; S, s) \rightarrow (Y, y, T, t)$, determined from the equation

$$M^{(s)}(u-\lambda)L^{(x)}(u) = L^{(y)}(u)M^{(t)}(u-\lambda) \quad (2.2)$$

and having the generating function $F_\lambda(y, t; x, s)$

$$X = \frac{\partial F_\lambda}{\partial x}, \quad Y = -\frac{\partial F_\lambda}{\partial y}, \quad S = \frac{\partial F_\lambda}{\partial s}, \quad T = -\frac{\partial F_\lambda}{\partial t} \quad (2.3)$$

(for simplicity, we omit the indices i in X_i, x_i etc.)

Let us impose now the constraint

$$t = s, \quad T = S. \quad (2.4)$$

Suppose that one can resolve the equations (2.3) and (2.4) with respect to $s = t$ and express X and Y as functions of (x, y) .

Proposition. The resulting transformation $B_\lambda(X, x) \rightarrow (Y, y)$ is canonical and is given by the generating function $\Phi_\lambda(x, y) = F_\lambda(y, s(x, y); x, s(x, y))$, such that

$$X = \frac{\partial \Phi_\lambda}{\partial x}, \quad Y = -\frac{\partial \Phi_\lambda}{\partial y}. \quad (2.5)$$

Proof. Let $|_{st}$ mean the restriction on the constraint manifold $s = t = s(x, y)$. The proof consists of two lines:

$$\begin{aligned} X &= \frac{\partial \Phi_\lambda}{\partial x} = \frac{\partial F_\lambda}{\partial x} \Big|_{st} + \frac{\partial F_\lambda}{\partial s} \Big|_{st} \frac{\partial s}{\partial x} + \frac{\partial F_\lambda}{\partial t} \Big|_{st} \frac{\partial t}{\partial x} \\ &= \frac{\partial F_\lambda}{\partial x} \Big|_{st} + \frac{\partial s}{\partial x} \left(\frac{\partial F_\lambda}{\partial s} + \frac{\partial F_\lambda}{\partial t} \right) \Big|_{st}. \end{aligned} \quad (2.6)$$

We observe now that

$$\left(\frac{\partial F_\lambda}{\partial s} + \frac{\partial F_\lambda}{\partial t} \right) \Big|_{st} = 0 \quad (2.7)$$

due to $S = T$, and, consequently, $X = \partial \Phi_\lambda / \partial x$. Similarly, one proves $Y = -\partial \Phi_\lambda / \partial y$.

To conclude, we notice that the constraint (2.4) implies the identity $M^{(s)}(u - \lambda) = M^{(t)}(u - \lambda)$ and, consequently, the equality (2.2) turns into the equality (1.4) ensuring thus that the transformation $\mathcal{B}_\lambda$ preserves the spectrum of $L(u)$.

3. Modification for the chain

In many applications the Lax matrix $L(u)$ is a monodromy matrix [2] factorized into the product of local Lax matrices $\ell_i(u)$

$$L(u) = \ell_N(u) \dots \ell_2(u) \ell_1(u) \quad (3.1)$$

having the same Poisson brackets (1.2) as $L(u)$

$$\{\ell_i^1(u), \ell_j^2(v)\} = [r_{12}(u - v), \ell_i^1(u) \ell_j^2(v)] \delta_{ij}. \quad (3.2)$$

The similarity transformation (1.4) is replaced now with a gauge transformation:

$$M_i(u - \lambda) \ell_i^{(x)}(u) = \ell_i^{(y)}(u) M_{i-1}(u - \lambda) \quad (3.3)$$

which ensures the preservation of the spectral invariants of $L(u)$ due to $M_N(u - \lambda) L^{(x)}(u) = L^{(y)}(u) M_N(u - \lambda)$.

The modification of the reduction procedure described in the previous section is quite straightforward. Supposing that $\ell_i(u)$ and $M_i(u)$ depend on local canonical variables

$$\ell_i^{(x)}(u) \equiv \ell_i(u; X_i, x_i), \quad \ell_i^{(y)}(u) \equiv \ell_i(u; Y_i, y_i), \quad (3.4a)$$

$$M_i^{(s)}(u) \equiv M(u; S_i, s_i), \quad M_i^{(t)}(u) \equiv M(u; T_i, t_i), \quad (3.4b)$$

we define first the local canonical transformations $\mathcal{R}_\lambda^{(i)} : (X_i, x_i; S_i, s_i) \rightarrow (Y_i, y_i, T_i, t_i)$ from the equations

$$M_i^{(s)}(u - \lambda) \ell_i^{(x)}(u) = \ell_i^{(y)}(u) M_i^{(t)}(u - \lambda). \quad (3.5)$$

Let the corresponding generating functions are $f_\lambda^{(i)}(y_i, t_i; x_i, s_i)$. Consider the direct product of N phase spaces $(X_i, x_i; S_i, s_i)$ and, respectively, $(Y_i, y_i; T_i, t_i)$. The generating function

$$F_\lambda := \sum_{i=1}^N f_\lambda^{(i)}(y_i, t_i; x_i, s_i) \quad (3.6)$$

determines then the direct product $\mathcal{R}_\lambda$ of the local canonical transformations $\mathcal{R}_\lambda^{(i)}$.

Let us now impose the constraint

$$t_i = s_{i-1}, \quad T_i = S_{i-1} \quad (3.7)$$

assuming periodicity $i + N \equiv i$. The proof of the canonicity of the resulting transformation $\mathcal{B}_\lambda : (X, x) \rightarrow (Y, y)$ parallels the proof given in the previous section. It remains to notice that after imposing the constraint (3.7) we have $M_i^{(s)}(u - \lambda) = M_i^{(t)}(u - \lambda)$ and obtain the equality (3.3).

4. Classical DST -model

We shall illustrate the general scheme described above on the simple example of the so-called DST -model — a degeneration of the Heisenberg spin chain [2]. In the notation of the previous section, the model is characterized by the local Lax matrices

$$\ell_i^{(x)}(u) = \begin{pmatrix} u - c_i - X_i x_i & x_i \\ -X_i & 1 \end{pmatrix} \quad (4.1)$$

whose Poisson brackets (3.2) are governed by the standard $SL(2)$ -invariant r -matrix [2]

$$r_{12}(u) = -\mathcal{P}_{12} u^{-1}, \quad (4.2)$$

where $\mathcal{P}_{12}$ is the permutation operator in $\mathbb{C}^2 \otimes \mathbb{C}^2$.

Let us choose for $M_i^{(s)}(u)$ another realization of the same Poisson brackets

$$M_i^{(s)}(u) = \begin{pmatrix} 1 & -S_i \\ s_i & u - S_i s_i \end{pmatrix}. \quad (4.3)$$

Note that our choice (4.3) of ansatz for $M_i^{(s)}(u)$ is different from (1.5) used in our previous paper [1]. In a sense, (4.3) is the simplest possible choice since $\det M(u - \lambda) = u - \lambda$ has only one pole instead of two as for (1.5).

From (3.5) one easily finds the local canonical transformation $\mathcal{R}_\lambda$ which is given by the equations

$$X = s, \quad Y = t + \frac{(c - \lambda)s}{1 - sy}, \quad (4.4a)$$

$$T = y, \quad S = x - \frac{(c - \lambda)y}{1 - sy}, \quad (4.4b)$$

or, equivalently, by the generating function

$$f_\lambda(yt \mid xs) = xs - yt + (c - \lambda) \ln(1 - sy). \quad (4.5)$$

Following the general scheme, we impose now the constraint (3.7). Using (4.4b) and resolving the equations $S_i = T_i$ with respect to s_i one obtains

$$s_i = \frac{1}{y_i} + \frac{c_i - \lambda}{y_{i+1} - x_i} \quad (4.6)$$

and, after substitution to (4.4a),

$$X_i = \frac{1}{y_i} + \frac{c_i - \lambda}{y_{i+1} - x_i}, \quad (4.7a)$$

$$Y_i = \frac{1}{y_{i-1}} + \frac{x_i - y_{i+1}}{y_i^2} + \frac{c_{i-1} - \lambda}{y_i - x_{i-1}} - \frac{c_i - \lambda}{y_i}. \quad (4.7b)$$

It is easy to verify that the equations (4.7) describe a canonical transformation whose generating function is

$$\Phi_\lambda(y \mid x) = \sum_i \frac{x_i - y_{i+1}}{y_i} + (c_i - \lambda) \ln \frac{y_i}{y_{i+1} - x_i}. \quad (4.8)$$

As expected from the general theory, the generating function $\Phi_\lambda(y \mid x)$ can also be obtained, up to a constant term, from F_λ corresponding to (4.5) by the substitution of s_i and t_i .

From the general proof given in sections 2 and 3 it follows that taking as ansatz for $M_i(u)$ any realization of the Poisson brackets (2.1) we will obtain some Bäcklund transformation. In particular, we can take for $M_i(u)$ a product of elementary matrices like (4.3). The corresponding Bäcklund transformation will be obviously the composition of the elementary Bäcklund transformations corresponding to (4.3). For instance, in such a way one can recover the matrix (1.5) and the corresponding two-parametric Bäcklund transformation $\mathcal{B}_{\lambda_1 \lambda_2}$ from [1].

5 Discussion

Our exposition was very sketchy, and we are planning to publish a more detailed study of our new technique elsewhere. Here we remark only that applicability of our construction goes far beyond the integrable systems connected to the $SL(2)$ -invariant r -matrice (4.2). The proof given in section 2 works for any solution to the classical Yang-Baxter equation provided the existence of the local canonical transformation $\mathcal{R}_\lambda$ and the conditions of degeneracy. The work on the application of the reduction technique to other integrable systems is in progress.

Another important remark concerns the connection with the quantum case. As noticed first in [4], see also [3], the quantum analog of Bäcklund transformation is the famous Baxter's Q -operator [5]. It can be shown that our construction of $\mathcal{B}_\lambda$ is the exact classical analog of the quantum construction of $Q(u)$ as a trace of a quantum monodromy matrix. Using this analogy one can, for example, construct the Q -operator for the quantum DST model [6].

Acknowledgements

I am grateful to V. B. Kuznetsov and F. Nijhoff for their interest in the work and useful discussions. This work was started during my stay at the Department of Mathematics, the University of Leeds and benefited from the financial support of EPSRC.

References

- [1] E.K. Sklyanin. *Canonicity of Bäcklund transformation: r-matrix approach. I*, solv-int/9903016.
- [2] L.D. Faddeev and L.A. Takhtajan, *Hamiltonian Methods in the Theory of Solitons*, Berlin: Springer (1987).
- [3] V.B. Kuznetsov and E.K. Sklyanin, *On Bäcklund transformations for many-body systems*, J. Phys. A: Math. Gen. **31** (1998) 2241–2251.
- [4] Pasquier V and Gaudin M 1992 The periodic Toda chain and a matrix generalization of the Bessel function recursion relations *J. Phys. A: Math. Gen.* **25** 5243–5252.
- [5] Baxter R I 1982 *Exactly Solved Models in Statistical Mechanics* (London: Academic Press, London) chapters 9–10.
- [6] E.K. Sklyanin, V.B. Kuznetsov and M. Salerno, *Quantum Bäcklund transformation for DST dimer model*, in preparation.